

Chlorpheniramine Maleate

COMPOSITION

Antista® Syrup : Each 5 ml syrup contains Chlorpheniramine Maleate BP 2 mg.

PHARMACOLOGY

Antista[®] (Chlorpheniramine Maleate) is an alkylamine antihistamine. It is one of the most potent H₁ blocking agents and is generally effective in relatively low doses. Antista is not so prone to produce drowsiness, readily absorbed from the gastro-intestinal tract, metabolised in the liver and excreted usually mainly as metabolised in the urine.

INDICATION

Antista® is mainly indicated in allergic conditions including urticaria, sensitivity reactions, angioneurotic oedema, seasonal hay fever, vasomotor rhinitis, cough, common cold, motion sickness

DOSAGE & ADMINISTRATION

Adults : 4 mg 3-4 times daily.

Children: Upto 1 (one) year: 1 mg twice daily. 1-5 years: 1 mg 3 - 4 times daily.

6-12 years : 2 mg 3-4 times daily.

or as directed by the physician.

SIDE EFFECT

Drowsiness, dizziness, headache, psychomotor impairment, urinary retention, dry mouth, blurred vision and gastro-intestinal disturbances, paradoxical stimulation may rarely occur, especially in high dosage or in children.

CONTRAINDICATION & PRECAUTION

There is no definite contraindication to therapy. It should be used with caution in epilepsy, prostatic hypertrophy, glaucoma and hepatic disease.

Warning: The ability to drive or operate machinery may be impaired.

STORAGE

Store below 30°C . Protect from light. Keep out of reach of children.

HOW SUPPLIED

Antista® Syrup: Bottle containing 60 ml / 100 ml syrup.

Manufactured by

