

# Espa-Vet<sup>®</sup> Injection

Drotaverine Hydrochloride INN

**Composition:**

Each ml injection contains Drotaverine Hydrochloride INN 20 mg.

**Indication:**

As antispastic in adjuvant treatment of digestive conditions (abdominal colic, gastric or duodenal ulcer, gastritis, enteritis, colitis, pancreatitis, spastic constipation, meteorism, spasms of cardia and pyloric sphincter, spasms of the biliary smooth muscles, biliary lithiasis, cholecystitis); spasms of urinary tract smooth muscles (renal urethral lithiasis, pyelitis, cystitis); genital disorders (uterine tetanic contractions, severe pains, abortion imminence) in dog, cat, horse and cattle. The spasmolytic action of the product also causes pain killing up to removal.

**Dosage & Administration:**

The product is administered by parenteral route: subcutaneous (S.C.), intramuscular (I.M.) or intravenous (I.V.).

*In Horses and Cattle:* 160 mg–200 mg active substance (8-10 ml product) by I.M. or S.C. route. Repeat the dose 2–3 times/day if necessary. In case of acute abdominal colic, inoculate slowly 8 ml product by I.V. route. The dose may be repeated 2-3 times/day by I.M. or S.C. route

*In Dog:* 1-2 ml by I.M. or S.C. route 1–2 times/day.

*In Cat:* 0.5 ml–1 ml by I.M. or S.C. route 1–2 times/day.

**Contraindication:**

Severe hepatic, renal or cardiac failure.

**Precautions:**

This drug is not to be administered to horses that are to be slaughtered for use in food. The administration is carried out only by the veterinarian or under his supervision. If administering by intravenous route, inject the product very slowly. Administer during pregnancy and lactation only after previous assessment of the risk/benefit therapeutic ratio.

**Side effects:**

Arterial blood pressure may drop following too rapid intravenous injection.

**Use in Pregnancy and Lactation:**

There is no known case of teratogenicity in animal studies. However, the use of Drotaverine should be avoided during pregnancy and lactation.

**Drug Interaction:**

Levodopa (anti-perkinson effect may be decreased).

**Storage:**

Store below 25° C & protect from light. Keep out of reach of children.

**Pack Size:** 10 ml vial.

For Veterinary Use only.

Manufactured by:



**SQUARE**  
PHARMACEUTICALS LTD.  
AGROVET DIVISION