

Cozin-VetTM

Zinc, Copper & Organic Acids

Solution

Composition

Zinc Sulfate 15%, Copper Sulfate 8%, Lactic Acid 8% & Acetic Acid 10%.

Indication

Prevents nonspecific diarrhea in layer and broiler. Improves FCR in broiler & increases egg production in layer. Prevents retarded growth and ensures flock uniformity of broiler. Prevents cannibalism and poor feather development in poultry. Reduces growth of harmful bacteria and helps to increase beneficial bacteria in the gut.

Dosage & Administration

Poultry: 1 ml/2 Liter of drinking water for 5-7 days.
Or as advised by the registered veterinary doctor.

Drug interaction

Concomitant administration of Tetracycline & Zinc may decrease the absorption of both the Tetracycline & Zinc. Similarly, concomitant administration of Zinc & Quinolone may also decrease the absorption of both.

Side effects

It has no side effects at recommended dose.

Precaution

Avoid contact with skin and eyes. If it comes in contact with skin or eyes, immediately flush the affected area with plenty of water.

Withdrawal period

Nil.

Storage

Store below 30° C temperature. Protect from light. Keep out of reach of children.

Pack size

100 ml & 1 Liter bottle.

For veterinary use only.

Manufactured by



SQUARE
PHARMACEUTICALS LTD.
AGROVET DIVISION
BSCIC I/A, Pabna, Bangladesh