

Ectonil-Vet[®] Powder

Metrifonate USP

Composition

Each gm powder contains Metrifonate USP (Trichlorfon) 1 gm.

Indication

Biting and sucking lice, mange mites, fleas, flies, red avian mites (*Dermanyssus avium*, *Dermanyssus gallinae*), scaly leg mites (*Cnemidocoptes mutans*), skin larvae of *Hypoderma bovis*, *Hypoderma lineatum*, *Dermatobia hominis*, stomach bots (*Gasterophilus spp.*), tick and roundworms (Hump sore).

Dosage & Administration

Cattle, pig, horse, sheep and goat: Wash or spray treatment of the whole animal; repeat after 5 days. Concentration 0.15% solution (1.5 gm per liter of water). Oral administration: 50-75 mg/kg body weight.

Poultry: Carefully spray or dip the fowl for 1-2 minutes. Spray also floors, walls, ceiling, nests, windows, doors and the other sites where the parasites or flies gather. Concentration 1.5 gm per liter of water.

Contraindication

Patients with a history of hypersensitivity to Metrifonate USP (Trichlorfon).

Use in Pregnancy and Lactation

Ectonil-Vet[®] Powder has not been shown to produce reproductive or teratogenic effects. But, do not administer to lactating dairy cows.

Adverse effects

No adverse effects are expected if this product is used in accordance with the label. Some symptoms of overexposure that can arise are: blurred vision, weakness, giddiness, sweating, muscular twitching, spasms, miosis and respiratory paralysis. Can be absorbed through skin. May irritate the skin. May irritate the eyes as well.

Storage

Store below 30° C. Protect from light. Keep out of reach of children.

Pack size

5 X 10 gm.

For veterinary use only.

To be dispensed only by or on the prescription of a registered veterinary doctor.

Manufactured by:



SQUARE
PHARMACEUTICALS LTD.
AGROVET DIVISION
Salgaria, Pabna, Bangladesh