



Fungidal®

Miconazole Nitrate
Antifungal

COMPOSITION

Cream : Each gram of Fungidal® topical cream contains Miconazole Nitrate BP 20 mg.

PHARMACOLOGY

Fungidal® is a broad-spectrum antimycotic which offers a high antifungal activity against dermatophytes, yeasts and other phyco-Asco and Adelomycetes, with a potent antibacterial activity against Gram-positive bacilli and cocci.

Fungidal® proved to be markedly effective in secondary infected mycoses, which under other treatments were resistant or reappeared. Fungidal® cream does not stain skin or clothes.

INDICATION

Skin and nail infections due to dermatophytes, yeasts and other fungi such as: Tinea capitis, corporis, manuum, pedis, barbae, cruris, unguium or onychomycosis. Pityriasis versicolor, candidiasis of skin and nails, stomatitis angularis, otitis externa.

Since Fungidal® has an antibacterial effect on Gram-positive bacteria, it may be used in mycoses secondarily infected with such bacteria.

DOSAGE AND ADMINISTRATION

The dosage is same for all the ages.

For skin infections: Apply some cream to the lesions twice daily and rub it well with finger until it has fully penetrated the skin.

All lesions usually disappear after 2 to 5 weeks. Prolong treatment for some 10 days to prevent relapse.

For nail infections: Clip infected nail as shortly as possible. Apply some cream once daily to the infected nail and rub with your finger, cover nail with a non-perforated occlusive plastic bandage.

Also after loosening of the infected nail (from 2-3 weeks onwards) uninterrupted treatment should be continued until the growth of a new nail has set in and definite cure can be observed (usually after seven months or more).

CONTRAINDICATION AND PRECAUTION

No contraindication is known.

Miconazole Nitrate is intended for topical use. If accidental ingestion of large quantities of the product occurs, an appropriate method of gastric emptying may be used if considered desirable.

SIDE EFFECT

Topical application of Miconazole Nitrate has almost no side effect.

USE IN PREGNANCY AND LACTATION

Only small amounts of Miconazole nitrate are absorbed following local administration. However as with other imidazoles, Miconazole nitrate should be used with caution during pregnancy.

STORAGE CONDITION

Store away from direct heat. Keep out of reach of children.

HOW SUPPLIED

Fungidal® : Tube containing 10 gm cream.

