



# Panvit<sup>®</sup>

**Multivitamin Paediatric Drops**

## COMPOSITION

*Each 0.6 ml contains :*

Vitamin A	BP	3000 IU
Vitamin D	BP	600 IU
Vitamin B <sub>1</sub> (Thiamine hydrochloride)	BP	0.96 mg
Vitamin B <sub>2</sub> (Riboflavin 5- phosphate sodium)	BP	0.60 mg
Vitamin B <sub>6</sub> (Pyridoxine hydrochloride)	BP	0.60 mg
Nicotinamide	BP	6.0 mg
Calcium D-pantothenate	USP	3.0 mg
Ascorbic acid	USP	30 mg

in palatable flavoured base

## PHARMACOLOGY

Each component of Panvit<sup>®</sup> in logical therapeutic concentration essential for maintenance of good and sound health of infants and children. Vitamin A plays an essential role in the function of retina and is essential for the growth and differentiation of epithelial tissues. Vitamin D is supplemented for prevention and cure of nutritional and metabolic rickets and for treatment of hypoparathyroidism. Deficiencies of B vitamins include glossitis, stomatitis, cheilosis, polyneuritis, beri-beri, pellagra, and vascularisation of cornea. Vitamin B-complex forms a part of co-enzyme responsible for various metabolic processes and promotes appetite. Ascorbic acid is used for the treatment of Vitamin C deficiency, especially causing scurvy in infants. Panvit<sup>®</sup> is balanced combination of vitamins which replenishes vitamin deficiency in children and can maintain the proper growth and resistance of the body against diseases.

## INDICATION

For prevention and treatment of threatened, mild, latent subclinical multivitamin deficiency. To meet the increased demand during the growth and illness. Dietary supplement in malnourished infants and children. As an adjunct therapy with antibiotic, sulphonamide, and isoniazide.

## DOSAGE AND ADMINISTRATION

Panvit<sup>®</sup> may be dropped directly in to the mouth or mixed with cereal, milk, fruit juices or other food. Usually 10-20 drops daily in divided doses or as advised by the physicians.

VITAMINS

## Panvit®

---

### **CONTRAINDICATION AND PRECAUTION**

It is possible to develop symptoms of taking excessive intake of vitamin D if the individual receives vitamin D fortified nutrients such as milk, infant formulas/foods in addition to Panvit®. It is also contraindicated to patients hypersensitive to any of its components.

### **SIDE EFFECT**

Side effects have been reported with specific vitamins, but at level substantially higher than those in Panvit®. Usually large dose of Panvit® should not be given.

### **DRUG INTERACTION**

No hazardous drug interaction was reported.

### **USE IN PREGNANCY AND LACTATION**

Vitamins present in Panvit® are safe for pregnant and nursing mothers. However, Panvit® is formulated only for paediatric patients.

### **HOW SUPPLIED**

Panvit® Paediatric Drops: Each bottle contains 15 ml drops with dropper.

