



# Servin®

Dried Ferrous Sulphate, Folic acid & Vitamins  
**Heamatinic**

## COMPOSITION

Servin® capsule : Each Servin® capsule contains timed release formulation of - Dried Ferrous Sulphate BP 150 mg, Folic acid BP 0.5 mg, Thiamine Mononitrate USP 2 mg, Riboflavin USP 2 mg, Nicotinamide USP 10 mg, Pyridoxine Hydrochloride BP 1 mg and Ascorbic Acid USP 50 mg.

## INDICATIONS AND USAGE

Servin® is indicated for the treatment and prophylaxis of iron, folic acid, vitamin B-complex and vitamin C deficiency, especially during pregnancy and lactation.

## DOSAGE AND ADMINISTRATION

One capsule daily. In more severe cases, 2 capsules a day may be required or as directed by the physician.

## CONTRAINDICATION AND PRECAUTION

This product is contraindicated in patients with a known hypersensitivity to any of the ingredients.

Care should be taken in patients who may develop iron overload, such as those with hemochromatosis, haemolytic anaemia or red cell aplasia. Iron chelates with tetracycline and absorption may be impaired.

## SIDE EFFECT

Generally well tolerated. However, a few allergic reactions may be seen.

## USE IN PREGNANCY AND LACTATION

Recommended.

## DRUG INTERACTION

No such drug interactions have been reported.

## OVERDOSAGE

Accidental overdose of iron containing products is a leading cause of fatal-poisoning in children fewer than 6 years. Avoid higher doses in liver disease or hemochromatosis; excess iron can cause bloody diarrhoea, vomiting, acidosis, darkened stools and abdominal pain. Symptoms may clear in a few hours. Riboflavin is reported to be completely safe and no toxic symptoms have been reported so far. Higher doses of Nicotinamide

HAEMATINICS

HAEMATINICS

## Servin<sup>®</sup>

---

may cause vomiting and diarrhoea. Sensory neuropathy was observed in individuals consuming more than 200 mg Pyridoxine for very long periods. No cases of Folic acid overdose have been reported. Acute ingestion of Ascorbic acid, even of massive doses, is unlikely to cause significant effects.

### **STORAGE CONDITION**

Store at a cool and dry place, protected from light and moisture.

### **HOW SUPPLIED**

Servin<sup>®</sup> capsule : Box containing 10 x 10 capsules in blister strips.



HAEMATINICS

HAEMATINICS