

# Antista<sup>®</sup>

Chlorpheniramine maleate

## COMPOSITION

Antista<sup>®</sup> Syrup : Each 5ml syrup contains Chlorpheniramine maleate BP 2 mg.

## PHARMACOLOGY

Antista<sup>®</sup> (Chlorpheniramine maleate) is an alkylamine antihistamine. It is one of the most potent H<sub>1</sub> blocking agents and is generally effective in relatively low doses. Antista<sup>®</sup> is not so prone to produce drowsiness, readily absorbed from the gastro-intestinal tract, metabolised in the liver and excreted usually mainly as metabolised in the urine.

## INDICATION

Mainly in allergic conditions including urticaria, sensitivity reactions, angioneurotic oedema, seasonal hay fever, vasomotor rhinitis, cough, common cold, motion sickness.

## DOSAGE AND ADMINISTRATION

*Adults* : : 4 mg 3-4 times daily.

*Children* : Up to 1 ( one) year : 1 mg twice daily.

1-5 years : 1 mg 3-4 times daily.

6-12 years : 2 mg 3-4 times daily.

or as directed by the physician.

## SIDE EFFECT

Drowsiness, dizziness, headache, psychomotor impairment, urinary retention, dry mouth, blurred vision and gastro-intestinal disturbances, paradoxical stimulation may rarely occur, especially in high dosage or in children.

## CONTRAINDICATION AND PRECAUTION

There is no definite contraindication to therapy. It should be used with caution in epilepsy, prostatic hypertrophy, glaucoma and hepatic disease.

*Warning:* The ability to drive or operate machinery may be impaired.

## HOW SUPPLIED

Antista<sup>®</sup> Syrup : Bottle containing 100ml syrup.

*Manufactured by :*



**SQUARE**

**PHARMACEUTICALS LTD.**

**BANGLADESH**

® Registered Trade Mark.