

## COMPOSITION

Erian® Ointment: Each gram ointment contains

Hydrocortisone BP 5 mg + Framycetin sulphate

Erian® Suppository: Each Suppository contains

Hydrocortisone BP 5 mg + Framycetin sulphate

## PHARMACOLOGY

The local anaesthetic cinchocaine hydrochloride

sometimes encountered in strangulated haemorrhoids

whilst the corticosteroid (Hydrocortisone) acts as

anti-pruritic agent and by so doing eliminates

discharge. The broad spectrum antibiotic, Framycetin

infections which may already be present or arising

Esculin has a skin protective action.

## INDICATION

For the treatment of internal and external haemorrhoids

Haemorrhoids post-partum

Anal pruritus, peri-anal eczema, anal fissures

Post-haemorrhoidectomy application to relieve

## DOSAGE & ADMINISTRATION

Ointment: Apply the ointment in a small quantity to the

pruritic area, morning and evening and after

elasticity, dilatation of superficial blood vessels, and other changes are particularly likely to occur when there is absorption of topically applied corticosteroids under the following conditions: when large quantities are applied to areas of the body, or to damaged skin; when prolonged use is made when the occlusive dressing technique is applied; when the pituitary-adrenal axis with consequent suppression of adrenal effects are most likely to be severe in children; and when a state may be produced. Benign intracranial hypertension

## DRUG INTERACTION

Proper data is not available.

## USE IN PREGNANCY

Pregnancy: The safe use of topical corticosteroids in pregnancy is established. Therefore, during pregnancy, the use of these drugs on extended areas, in large amounts or for prolonged periods

## USE IN CHILDREN

Not recommended

## STORAGE

Ointment: Store at a cool and dry place protected from light.

Suppository: Store below 25°C (77°F).