

Erian[®]

Cinchocaine Hydrochloride + Hydrocortisone + Framycetin sulphate + Esculin

COMPOSITION

Erian[®] Ointment: Each gram ointment contains Cinchocaine Hydrochloride BP 5 mg + Hydrocortisone BP 5 mg + Framycetin sulphate BP 10 mg + Esculin 10 mg.

Erian[®] Suppository: Each Suppository contains Cinchocaine Hydrochloride BP 5 mg + Hydrocortisone BP 5 mg + Framycetin sulphate BP 10 mg + Esculin 10 mg.

PHARMACOLOGY

The local anaesthetic cinchocaine hydrochloride prevents or relieves the severe pain sometimes encountered in strangulated haemorrhoids, fissures and perianal haematomata, whilst the corticosteroid (Hydrocortisone) acts as a decongestant, anti-inflammatory and anti-pruritic agent and by so doing eliminates itching, inflammation and mucous discharge. The broad spectrum antibiotic, Framycetin sulphate, will eradicate most infections which may already be present or arise in lesions of the anorectal area and Esculin has a skin protective action.

INDICATION

For the treatment of internal and external haemorrhoids

Haemorrhoids post-partum

Anal pruritus, peri-anal eczema, anal fissures and proctitis

Post-haemorrhoidectomy application to relieve pain and discomfort

DOSAGE & ADMINISTRATION

Ointment: Apply the ointment in a small quantity with the finger, on the painful or pruritic area, morning and evening and after each stool. For deep application attach cannula/applicator (Supplied) to tube, insert to full extent and squeeze tube gently from lower end whilst withdrawing.

Suppository: A suppository is inserted morning and evening, and after each stool.

CONTRAINDICATION

Known hypersensitivity to any of the four ingredients. Corticosteroids have been shown to be teratogenic in animals following dermal application. As these agents are absorbed percutaneously, teratogenicity following topical application cannot be excluded. Therefore **Erian[®]** should not be used during pregnancy. Topical corticosteroid preparations are contra-indicated in the treatment of herpes simplex, vaccinia or varicella, or tuberculous infection of the anal region.

PRECAUTION

Discontinue use if sensitization occurs. Other specific measures against infections, allergy, and other causal factors must not be neglected. The possibility, however rare, that prolonged use of this preparation might produce systemic corticosteroid effects, should be borne in mind. Patients should be advised to inform subsequent physicians of the previous use of hydrocortisone.

SIDE-EFFECTS

Long-term continuous treatment with topical corticosteroids should be avoided as far as possible as this may cause atrophic changes in the skin leading to thinning, loss of elasticity, dilatation of superficial blood vessels, telangiectasia and ecchymoses. These changes are particularly likely to occur when occlusive dressings are used. Systemic absorption of topically applied corticosteroids may occur, particularly under the following conditions: when large quantities are used or when application is made to wide areas of the body, or to damaged skin; when potent topical corticosteroids are used, and when the occlusive dressing technique is applied. Depression of the hypothalamic-pituitary-adrenal axis with consequent suppression of the adrenal gland may occur. These effects are most likely to be severe in children. Growth may be retarded and a Cushingoid state may be produced. Benign intracranial hypertension has been rarely reported.

DRUG INTERACTION

Proper data is not available.

USE IN PREGNANCY

Pregnancy: The safe use of topical corticosteroids during pregnancy has not been fully established. Therefore, during pregnancy, they should not be used unnecessarily on extended areas, in large amounts or for prolonged periods of time.

USE IN CHILDREN

Not recommended

STORAGE

Ointment: Store at a cool and dry place protected from light. Do not freeze.

Suppository: Store below 25° C.

HOW SUPPLIED

Erian[®] Ointment: Each pack has a tube containing 15 gm of ointment with an Applicator

Erian[®] Suppository: Box containing 2x5's Suppositories in blister pack.

Manufactured by :



SQUARE

PHARMACEUTICALS LTD.
BANGLADESH

© Registered Trade Mark.