

Repres[®] Plus

Perindopril+Indapamide
ACEI– Diuretic combination

Presentation:

Repres[®] plus 2 Tablet: Each tablet contains Perindopril Erbumine BP 2 mg + Indapamide BP 0.625 mg.

Repres[®] plus 4 Tablet: Each tablet contains Perindopril Erbumine BP 4 mg + Indapamide BP 1.25 mg

DESCRIPTION

Repres[®] plus is a fixed combination of an ACE inhibitor and a thiazide-like diuretic, leads to additive synergy of the antihypertensive effects of the two constituents. Its pharmacological are derived from those of each of the two components taken separately. In addition to those due to the additive synergistic action of the two constituents, when combined, it shows beneficial effects on vascular endothelium, arteriolo-capillary microcirculation and the target organs of hypertension,

INDICATIONS AND USES

Essential hypertension.

DOSAGE & ADMINISTRATION

1 tablet daily, preferably to be taken in the morning and before a meal.

Elderly patients: Normal dosage.

Renal failure: Creatinine clearance, CrCl >30 ml/min: no dosage modification.

CrCl <30 ml/min: treatment contraindicated.

CONTRAINDICATIONS

Absolute: Known allergy to perindopril, indapamide or sulfonamides; history of Quincke's edema linked to previous ACE inhibitor therapy; severe renal failure; serious liver disorder; hypokalemia; pregnancy; lactation.

Relative: Combination therapy with lithium, potassium salts, potassium-sparing diuretics, and certain medicines which can cause heart rhythm disorders.

SIDE EFFECTS

Asthenia, dizziness, headache, mood swings and/or sleep disturbances, cramps, hypotension, allergic reactions, skin rashes, gastrointestinal disorders, dry cough, dry mouth, risk of dehydration in the elderly and in patients suffering from heart failure; changes in blood test results may occur.

PRECAUTIONS

Disorders of electrolyte balance, diabetes, gout, hypotension, or strict sodium-free diets, heart or renal failure, atherosclerosis, renal artery stenosis, elderly.

DRUG INTERACTION

Not recommended: Combinations with lithium, potassium-sparing diuretics, potassium (salt), antiarrhythmic drugs which cause torsade de pointes, anesthetic drugs, cytostatic or immunosuppressive agents.

USE IN PREGNANCY AND LACTATION

As this combination contains an ACE inhibitor, it should not be used during pregnancy. Animal studies have shown that, perindopril and its metabolites are excreted in breast milk during lactation, but there are no human data. It is not known whether indapamide is excreted in breast milk. It is therefore recommended that the drug should not be given to lactating women, as the possible effect on the newborn is unknown.

USE IN PEDIATRIC PATIENTS

The safety and effectiveness in pediatric patients have not been established.

STORAGE CONDITION


Store in a cool and dry place. Protect from light and moisture.

HOW SUPPLIED

Repres[®] plus 2 Tablet: Each box contains 3x10 tablets in blister pack.

Repres[®] plus 4 Tablet: Each box contains 2x10 tablets in blister pack.

® Registered Trade Mark



**SQUARE
PHARMACEUTICALS LTD.
BANGLADESH**