

Bicozin-I[®]

Iron, Vitamin-B Complex & Zinc Syrup

COMPOSITION

Bicozin-I[®] syrup: Each 5 ml of syrup contains Iron (III) Hydroxide Polymaltose Complex INN 200 mg equivalent to elemental Iron 50 mg, Thiamine Hydrochloride BP 5 mg, Riboflavin-5-Phosphate Sodium BP 2.74 mg equivalent to Riboflavin 2 mg, Pyridoxine Hydrochloride BP 2 mg, Nicotinamide BP 20 mg and Zinc Sulphate Heptahydrate BP 43.97 mg equivalent to elemental Zinc 10 mg.

PHARMACOLOGY

Bicozin-I[®] syrup is the preparation of Iron, Vitamin B-complex and Zinc. In the preparation, Iron is present as Iron (III) Hydroxide Polymaltose Complex, a noble Iron preparation, which contains non-ionic ferric iron and Polymaltose in a stable complex. This facilitates a controlled absorption of the Iron when it comes in contact with the mucosal cell surface. Being non-ionic, Iron (III) Hydroxide Polymaltose Complex is more stable than conventional Iron form.

INDICATION

Bicozin-I[®] syrup is indicated for the treatment and prevention of Iron, B-vitamins and Zinc deficiencies.

DOSAGE AND ADMINISTRATION

Adults: 5 ml-10 ml (1-2 teaspoonful) 3 times daily or as recommended by the physician.

Children: 5 ml (1 teaspoonful) 3 times daily or as recommended by the physician.

Infants: 0.33 ml/kg body weight daily or as recommended by the physician.

SIDE-EFFECT

Bicozin-I[®] syrup is generally well tolerated. However, a few side-effects of oral Iron preparations, including nausea, vomiting, constipation or diarrhoea may occur.

PRECAUTION

Caution should be taken in the conditions where is a risk of Iron overload, such as haemochromatosis, thalassemia, haemosiderosis or haemolytic anaemia.

USE IN PREGNANCY AND LACTATION

Recommended.

CONTRAINDICATION

Bicozin-I[®] syrup is contraindicated in patients with a known hypersensitivity to any of the ingredients of this product.

DRUG INTERACTION

No interactions have been observed. Since, the Iron is complex bound, ionic interaction with foodstuff components (phytates, oxalates, tannin, etc.) and concomitant administrations of medicaments (tetracyclines, antacids) are unlikely to occur.

OVERDOSE

In case of overdose, initially epigastric pain, pain, diarrhoea and vomiting can occur and may include metabolic acidosis, convulsions and coma after apparent recovery. Should seek emergency medical attention in case of overdose. Initially an emetic should be given and then gastric lavage and general supportive measures should be employed.

STORAGE

Store in a cool and dry place, away from light. Keep out of reach of children.

HOW SUPPLIED

Bicozin-I[®] 100 ml syrup: Each pack has a PET bottle containing 100 ml syrup and a measuring spoon.

Bicozin-I[®] 200 ml syrup: Each pack has a PET bottle containing 200 ml syrup and a measuring spoon.

Manufactured by :



SQUARE

PHARMACEUTICALS LTD.
BANGLADESH

© Registered Trade Mark.