

Ziliron®

Iron (III) Hydroxide Polymaltose Complex, Folic Acid & Zinc Sulfate Monohydrate

COMPOSITION

Each film coated tablet contains Iron (III) Hydroxide Polymaltose Complex INN 188 mg equivalent to 47 mg elemental Iron, Folic Acid BP 0.50 mg and Zinc Sulfate Monohydrate USP 61.80 mg equivalent to 22.50 mg elemental Zinc.

PHARMACOLOGY

Ziliron® tablet is the preparation of Iron, Folic Acid and Zinc. In the preparation, Iron is present as Iron Polymaltose Complex, a novel iron preparation, which contains non-ionic ferric iron and Polymaltose in a stable complex. This facilitates a controlled absorption of the Iron when it comes in contact with the mucosal cell surface. Being non-ionic, Iron Polymaltose Complex is more stable than conventional Iron form. Folic acid helps in the proper development of the baby. Zinc keeps enzymes working and helps metabolize proteins.

INDICATION

Prevention and treatment of Iron, Folic Acid and Zinc deficiencies.

DOSAGE AND ADMINISTRATION

One tablet daily or two tablets may be required a day in severe cases or as directed by the physician.

SIDE-EFFECT

Ziliron® tablet is well tolerated. Occasionally gastrointestinal irritation such as sensation of repletion, pressure in the epigastric region, nausea, constipation or diarrhoea may occur.

PRECAUTION

A dark coloration of the stool may occur which is without clinical significance.

CONTRADICTION

Ziliron® tablet is contraindicated in patients with a known hypersensitivity to any of the ingredients of this product.

USE IN PREGNANCY

Use of any drug during the first trimester of pregnancy should be avoided if possible. Thus administration of Iron the first trimester requires definite evidence of Iron deficiency. Prophylaxis of Iron deficiency where inadequate diet calls for supplementary Zinc and Folic Acid is justified during the remainder of pregnancy.

DRUG INTERACTION

No interactions have been observed. Since, the iron is complex bound, ionic interaction with foodstuff components (phytates, oxalates, tannin, etc.) and concomitant administrations of medicaments (tetracyclines, antacids) are unlikely to occur.

OVERDOSE

No toxication or iron overload has been reported in case of overdose.

STORAGE

Store at a cool & dry place, protected from light & moisture. Keep out of reach of children.

HOW SUPPLIED

Ziliron® Tablet : Box containing 6X10's tablets in blister pack.

Manufactured by :



SQUARE
PHARMACEUTICALS LTD.
BANGLADESH



® Registered Trade Mark.