

Sultolin[®]

Salbutamol sulphate BP

COMPOSITION

Sultolin[®] 4 Tablet: Each tablet contains Salbutamol sulphate BP equivalent to Salbutamol 4 mg.

INDICATION

Sultolin[®] is indicated in bronchial asthma, status asthmaticus and allied conditions concerned with bronchospasm.

DOSAGE AND ADMINISTRATION

Adults: 2-4 mg three to four times daily.

Some patients require up to 8 mg. Elderly patient should be given the lower doses initially.

Children 2-6 years: 1 to 2 mg three to four times daily or as prescribed by the physician.

CONTRAINDICATION

Hypersensitivity to Salbutamol contraindicates the use of it.

PRECAUTION

Salbutamol should be used with caution in patients with hyperthyroidism, diabetes mellitus, severe cardiovascular disorders or hypertension. In asthmatic patients whose condition deteriorates despite of Salbutamol therapy the dosage should not merely be increased, but alternative or additional therapy including corticosteroids should be instituted promptly.

SIDE EFFECT

These are usually trivial, consisting mainly of tachycardia and tremor. Headache and peripheral vasodilatation may occur in large doses.

STORAGE

Store in a cool and dry place away from heat and moisture.

HOW SUPPLIED

Sultolin[®] 4 Tablet: Box containing 200 tablets.