

Rislon®

Betahistine Mesilate BP 6 mg

Rislon® dilates precapillary sphincters, increasing the blood flow in the inner ear. It controls the permeability of capillaries in the inner ear, thereby removes endolymphatic hydrops. It also improves cerebral circulation, increasing blood flow in the internal carotid artery. Thus, **Rislon®** is clinically useful for the relief of vertigo and dizziness.

COMPOSITION: Each tablet of **Rislon®** contains Betahistine Mesilate BP 6 mg.

INDICATIONS: Vertigo and dizziness associated with the following diseases: Meniere's disease, Meniere's syndrome, Peripheral vertigo.

MODE OF ACTION: Betahistine Mesilate increases microcirculation blood flow in the inner ear, and removes endolymphatic hydrops of the inner ear. It also significantly increases the blood flow of the internal carotid artery.

CLINICAL ADVANTAGES: For discomfort accompanying cerebral disturbances: **Rislon®** exhibits a marked healing effect on giddiness experienced upon standing, the sensation of instability, and symptoms of dizziness. The heavy feeling in head and the accompanying discomforts can also be successfully subdued. For dizziness and the sensation of dizziness due to disturbances in the inner ear: **Rislon®** improves microcirculation of the inner ear and removes endolymph hydrops, thereby alleviating dizziness in Meniere's disease and Meniere's syndrome. Tinnitus, the sensation of ear obstructions, nausea and vomiting are also effectively allayed. For the prevention of dizzy spells and other residual symptoms: Along with inhibiting the symptoms of dizziness and removing residual symptoms after attack, continued administration of **Rislon®** also prevents the recurrence of dizzy spells. As the primary in prescribing for dizziness: By the combined use with antihypertensive agents, therapeutic agents for arteriosclerosis, vitamins and tranquilizers, accompanying symptoms can also be brought into remission and better therapeutic effects obtained.


DOSAGE AND ADMINISTRATION: Usually for adults, administer orally 1 to 2 tablets three times per day after meals. The dose may be adjusted according to the age of patient and severity of symptoms.

PRECAUTIONS: Careful Administration: Patients with a history of digestive ulcer or an active digestive ulcer, Patients with bronchial asthma, Patients with pheochromocytoma. Adverse Reactions: Digestive – Nausea or vomiting may rarely occur. Hypersensitivity– Hypersensitivity reactions, such as skin rash, may rarely occur. Use during Pregnancy: Safety of **Rislon®** during pregnancy has not been established. This drug should be administered to pregnant patients or women suspected of being pregnant, only if the expected therapeutic benefit is thought to outweigh any possible risk

STORAGE AND HANDLING: Cautions: Use only pursuant to the prescription or under supervision of doctors, etc. Storage: **Rislon®** Tablets should be stored at room temperature and stored in a moisture–proof container after unsealing.

HOW SUPPLIED: Each box contains 10 x10 Tablets in blister packing.

Manufactured by:



SQUARE
PHARMACEUTICALS LTD.
BANGLADESH

® Registered Trade Mark