

Climycin™

Clindamycin

Active Ingredient

Each capsule contains 300 mg Clindamycin as Clindamycin HCl.

Indication

Climycin has been shown to be Effective in the treatment of the following infections when caused by susceptible anaerobic bacteria or susceptible strains of gram positive bacteria such as streptococci, staphylococci & pneumococci: Upper respiratory tract infections, Lower respiratory tract infections, Skin & soft tissue infections, Bone & joint infections, Pelvic infections, Intra-abdominal infections, Septicemia & endocarditis, Dental infections. As an alternative therapy when used in combination with quinine or amodiaquine for the treatment of multi-drug resistant Plasmodium falciparum infection.

Dosage & Administration

Adults: Serious Infections: 150 mg - 300 mg every six hours. More severe infections: 300 mg - 450 mg every six hours.

Pediatric Patients: Serious Infections:

8 - 16 mg/kg/day divided into three or four equal doses.

More severe infections: 16 - 20 mg/kg/day divided into three or four equal doses.

Contraindication & Precaution

Clindamycin is contraindicated in patients previously found to be sensitive to Clindamycin or lincomycin or any of the ingredients of this medicine.

Side Effect

The adverse effects have been reported with the use of Clindamycin are abdominal pain, oesophagitis & oesophageal ulcer, nausea, vomiting & diarrhoea, pruritus, skin rashes, urticaria.

Use in Pregnancy & Lactation

Pregnancy: Pregnancy Category B: Clindamycin should be used in pregnancy only if clearly needed.

Lactation : Clindamycin has been reported to appear in breast milk. Therefore, it is not recommended for nursing mothers if not clearly needed.

Use in Children

When Clindamycin is administered to newborns & infants (birth to 16 years) appropriate monitoring of organ system functions is desirable.

Preparation

Each box contains 30 capsules in Alu-PVC blister pack.

Manufactured by



SQUARE

**PHARMACEUTICALS LTD.
BANGLADESH**