Dermasol Plus[™] Ointment

Clobetasol Propionate BP 0.05% & Salicylic Acid BP 3%

Composition

Dermasol Plus™ Ointment: Each gram ointment contains Clobetasol Propionate BP 0.5 mg and Salicylic Acid BP 30 mg.

Pharmacology

Clobetasol Propionate, a synthetic corticosteroid and Salicylic Acid, a keratolytic agent for topical dermatologic use. Clobetasol is an analog of prednisolone having a high degree of glucocorticoid activity and a slight degree of mineralocorticoid activity. Salicylic Acid is a keratolytic and antiseptic agent.

Indication

Dermasol PlusTM Ointment is indicated for the relief of the inflammatory manifestations of hyperkeratotic and dry corticosteroid-responsive dermatitis such as psoriasis, chronic atopic dermatitis, neurodermatitis (lichen simplex chronicus), lichen planus, eczema (including nummular eczema, hand eczema, eczematous dermatitis), dyshidrosis (pompholyx), seborrheic dermatitis of the scalp, ichthyosis vulgaris and other ichthyotic conditions.

Dosage and Administration

Apply a thin layer of **Dermasol Plus**[™] Ointment to the affected skin areas twice daily and rub gently and completely.

Contraindication

This preparation is contraindicated in patients with a history of sensitivity to any components of the preparation.

Precaution

It should not be used on the face, groin or axillae. This ointment is not for ophthalmic use. If irritation or sensitization develops with the use of this ointment, treatment should be discontinued.

Side Effect

Burning sensation, irritation, itching, stinging sensation, erythema & telangiectasia.

Drug Interaction

It is not known whether it has any interaction with other topically applied drug.

Pregnancy & Lactation

Since safety of topical corticosteroid use in pregnant women has not been established, drugs of this class should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. Since it is not known whether topical administration of corticosteroids can result in sufficient systemic absorption to produce detectable quantities in breast milk, a decision should be made to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

Pediatric Use

The safety and effectiveness of this product have not been established in pediatric patients. Use 12 years of age is not recommended.

Storage

Store $\bar{i}n$ a dry and cool place below 30° C, protect from light. Do not refrigerate. Keep out of reach of children. Only use by the prescription of a registered physician.

How Supplied

Dermasol Plus™ Ointment: Each pack has an aluminun tube contains 30 gm Ointment.

Manufactured by

