

Antihista-Vet[®] Injection

Pheniramine Maleate BP

Composition:

Each ml injectable solution contains Pheniramine Maleate BP 22.75 mg

Indication:

Itching at different areas of body, drug allergy, hay fever, urticaria, inflammation, constipation, rhinitis, pulmonary oedema, local anaesthesia, bronchospasm, puerperal toxemia, inflammation of stomach and uterus.

Dosage and Administration:

Antihista-Vet[®] injection is administered by IM & IV route. Onset of action is quick following IV administration and acts within 10-20 minutes after IM administration. The drug can be administered 2-3 times daily, depending on the severity of the disease.

Cattle, Buffalo & Horse : 5-10 ml per day.

Goat & Sheep : 0.5-1 ml per day.

Or as directed by the registered veterinary physician.

Contraindication:

The drug is contraindicated in animal hypersensitive to active ingredient.

Side Effect:

In therapeutic doses, slight insomnia, drowsiness may develop & may feel fatigue. Injection may be irritant and transient hypotension or stimulation of the CNS.

Use in Pregnancy & Lactation:

Safety for the use of Antihista-Vet[®] injection during pregnancy has not been established. Antihista-Vet[®] injection should only be used when clearly needed and potential benefit outweighs the potential unknown risks to the fetus. Use during lactation is not recommended.

Storage:

Protect from light, store in a cool and dry place. Keep out of reach of children.

Pack Size:

10X10 ml & 100 ml.