

Cerevas

Vinpocetine

COMPOSITION

Cerevas® Tablet : Each tablet contains Vinpocetine INN 5 mg.

INDICATIONS

Acute Cerebro-Vascular Accidents (Strokes): Ischemic strokes due to cerebral thrombosis, cerebral embolism, acute circulatory disorder, hypertensive crisis; the acute cardiovascular disorders, ischemic neurological deficit (RIND) complete stroke (CS), multiinfarct dementia, cerebral arteriosclerosis, hypertensive encephalopathy, post-apoplectic conditions with the background of hemorrhagic strokes etc.

Geriatrics

For relief of psychosomatic symptoms in the elderly due to cerebral insufficiency e.g. forgetfulness, memory disturbances, slow thinking, lack of concentration, dizziness, mood instability, aphasia, sleep disturbances, vasovegetative symptoms of menopausal syndrome etc.

Ophthalmology

Vascular disorder of the choroid and retina due to arteriosclerosis. Vasospasm, macula degenerations, arterial or venous thrombosis or embolism or glaucoma secondary to above mentioned disorders.

Otology

For the treatment of impaired hearing of vascular toxic (iatrogenic) origin presbycusis. Meniere's disease, cochleovestibular neuritis, tinnitus and dizziness of labyrinth origin.

DOSAGE

1-2 tablets thrice daily, the maintenance dose is one tablet thrice daily.

CONTRA-INDICATIONS

It is contra-indicated in pregnancy.

PRECAUTIONS

In the acute case of above mentioned neurological, ophthalmological and otological disorders until the improvement of symptoms parenteral treatment is recommended followed by oral treatment. In chronic cases oral therapy should be applied.

SIDE EFFECTS

Transient hypotension, tachycardia may occur.

DRUG INTERACTIONS

Interactions of the tablet have not been reported so far therefore, it can be applied in combinations.

STORAGE

Store in a cool and dry place, protected from light and moisture.

HOW SUPPLIED

Cerevas® Tablet : Each box contains 5x10's tablets in blister packing.