

Ciprocin-Vet[®] Solution

Ciprofloxacin USP

Composition:

Each ml Solution contains Ciprofloxacin USP 100 mg as hydrochloride.

Indication:

Ciprocin-Vet[®] Solution is effective against several diseases as C.R.D, C.C.R.D, Mycoplasmosis, Salmonellosis, Colibacillosis, Infectious Coryza, Pasteurellosis. Streptococcosis, Staphylococcosis etc. and may be used in viral disease like Gumboro, Newcastle disease etc. followed by secondary bacterial infections.

Mechanism of Action:

Ciprofloxacin functions by inhibiting DNA-gyrase enzyme necessary to separate bacterial DNA, thereby inhibiting cell division.

Dosage and Administration:

Poultry:

Treatment: 1 ml per 1-2 litre of drinking water for 3-5 days.

Prevention: 1 ml per 4-8 liter of drinking water for 3 days.

or as directed by the Veterinarian.

Side Effects:

Central nervous system (CNS) toxicity; gastrointestinal reactions; hepatotoxicity; hypersensitivity reactions; photosensitivity; phototoxicity.

Drug Interaction:

Ciprofloxacin administered together with theophylline can lead to elevated blood levels of theophylline. Ciprofloxacin can enhance the action of the anticoagulant warfarin and increase the risk of bleeding.

Contraindication:

It should not be given to patients with a history of hypersensitivity to Ciprofloxacin or to other quinolones.

Precaution:

It should be used with caution in patients with suspected or known CNS disorders.

Storage:

Store below 30° C in a cool and dry place. Protect from light. Do not refrigerate. Keep out of reach of children.

Pack Size:

100 ml, 500 ml.