

Glysup™

Glycerin BP

COMPOSITION

Glysup™ 1.15 Suppository: Each suppository contains Glycerin BP 1.15 gm.

Glysup™ 2.30 Suppository : Each suppository contains Glycerin BP 2.30 gm.

PHARMACOLOGY

Glycerin is a hyperosmotic laxative, given rectally, which usually produces a bowel movement within 15 minutes to 1 hour. Hyperosmotic laxatives encourage bowel movements by drawing water into the bowel from surrounding tissues. This produces a softer stool mass and increased bowel action. These products are used for fast, predictable relief of occasional constipation.

INDICATION

For the relief of occasional constipation.

DOSAGE AND ADMINISTRATION

Children under 2 years: Consult a physician.

Children (2 to 6 years): only 1 **Glysup™** 1.15 suppository per 24 hours or as directed by a physician.

Adults and Children (From 6 years): only 1 **Glysup™** 2.30 suppository per 24 hours or as directed by a physician

Insert suppository well up into rectum. Suppository need to melt completely to produce laxative action.

CONTRAINDICATION

Sensitivity to the ingredients. Do not use unless the patient to be treated is, in fact, constipated.

SIDE EFFECTS

Glycerin when used rectally may cause rectal discomfort or a burning sensation.

USE IN PREGNANCY

Pregnancy category C. There are no controlled data in human pregnancy.

STORAGE

Store below 25°C. Protect from moisture.

HOW SUPPLIED

Glysup™ 1.15 Suppository: Box containing 30 suppositories in blister pack.

Glysup™ 2.30 Suppository: Box containing 20 suppositories in blister pack.

Manufactured by:

Manufactured by :



SQUARE

PHARMACEUTICALS LTD.
BANGLADESH

TM - Trade Mark

Revision No.00