

# Norvis<sup>®</sup> Tablet / Injection

Tiemonium methylsulphate INN

---

## Norvis<sup>®</sup> Tablet

**Composition:** Each film coated Tablet contains 50 mg Tiemonium methylsulphate INN.

**Pharmacology:** Tiemonium are quaternary ammonium antimuscarinic agent with peripheral effect similar to those of atropine. Tiemonium strengthens calcium bonding with phospholipids and proteins thus stabilizing the cell membrane of the GI tract.

**Indication and usage:** Tiemonium methylsulphate is an antispasmodic drug. It reduces muscle spasms of the intestine, biliary system, uterus & urinary bladder. It is indicated for the pain in gastrointestinal & biliary disease in the urology and gynecology such as gastroenteritis, diarrhea, dysentery, biliary colic, enterocolitis, cholecystitis, colonopathies, mild cystitis, & spasmodic dysmenorrhoea.

**Dosage and administration:**

Recommended oral dose of Tiemonium methylsulphate tablet is 2-6 tablets (100-300 mg) daily in divided dosage as required.

**Precaution:**

**Drug Interactions:** Tiemonium methylsulphate tablet should not be used with other drugs without prior consult of a registered physician to avoid possible drug interaction.

**Pregnancy:** Tiemonium methylsulphate tablet may be used in pregnancy only if it is clearly needed by the assessment of risk benefit ratio.

**Nursing mother:** Tiemonium methylsulphate tablet may be used in lactating mother only if it is clearly needed by the assessment of risk benefit ratio.

**Paediatric use:** safety and effectiveness of Tiemonium methylsulphate tablet in paediatric patients have not been established.

**Geriatric use:** Efficacy and safety were maintained with increasing age.

**Side effects:**

Side effect with the use of Tiemonium methylsulphate tablet is very rare.

**Contraindication:**

Tiemonium methylsulphate tablet should not be used in Glucoma or where acute pain of eyeball with vision disturbance was found. Also this drug should not be used in patients with the disorder of prostate or urinary bladder.

**Overdose:**

There is not available data regarding the overdose of Tiemonium methylsulphate tablet.

**How Supplied:**

Each box contains 5 x 10's tablets in blister pack.

## Norvis<sup>®</sup> Injection

**Composition:** Each 2 ml ampoule contains 5 mg of Tiemonium methylsulphate INN.

**Pharmacology:** Tiemonium are quaternary ammonium antimuscarinic agent with peripheral effect similar to those of atropine. Tiemonium strengthens calcium bonding with phospholipids and proteins thus stabilizing the cell membrane of the GI tract.

**Indication and usage:** Tiemonium methylsulphate is an antispasmodic drug. It reduces muscle spasms of the intestine, biliary system, uterus & urinary bladder. It is indicated for the pain in gastrointestinal & biliary disease in the urology and gynecology such as gastroenteritis, diarrhea, dysentery, biliary colic, enterocolitis, cholecystitis, colonopathies, mild cystitis, & spasmodic dysmenorrhoea.

**Dosage and administration:**

Recommended injectable dose is one or two Tiemonium methylsulphate Injection three times daily, through Intravenous route slowly or Intramuscular route.

**Precaution:**

Drug Interactions: Tiemonium methylsulphate should not be used with other drugs without prior consult of a registered physician to avoid possible drug interaction.

Pregnancy: Tiemonium methylsulphate may be used in pregnancy only if it is clearly needed by the assessment of risk benefit ratio.

Nursing mother: Tiemonium methylsulphate may be used in lactating mother only if it is clearly needed by the assessment of risk benefit ratio.

Paediatric use: safety and effectiveness of Tiemonium methylsulphate in paediatric patients have not been established.

Geriatric use: Efficacy and safety were maintained with increasing age.

**Side effects:**

Side effect with the use of Tiemonium methylsulphate is very rare.

**Contraindication:**

Tiemonium methylsulphate should not be used in Glucoma or where acute pain of eyeball with vision disturbance was found. Also this drug should not be used in patients with the disorder of prostate or urinary bladder.

**Overdose:**

There is not available data regarding the overdose of Tiemonium methylsulphate.

**How Supplied:**

Each box contains 2 x 5's of 2ml ampoules in blister pack.