

Prithi[®] WS Powder

Vitamin B₁, B₂ & B₆

Composition:

Each 100 gm powder contains -

Vitamin B ₁	5 gm
Vitamin B ₂	1 gm
Vitamin B ₆	2 gm
Dextrose monohydrate q. s. to make	100 gm.

Pharmacology:

The members of the vitamin B group contained in Prithi[®] WS Powder are components of enzyme systems that regulate various stages of carbohydrate, fat and protein metabolism, each of the components playing a specific biological role. Deficiency of B vitamin causes Polyneuritis, Curled toe paralysis, inappetance, dermatitis, glossitis, stomatitis etc.

Prithi[®] WS Powder, Each member of the B-complex has a unique structure and performs unique functions in the poultry's body. These vitamins are vital for:

- Vitamin B₁ (Thiamin) helps the body to break down carbohydrates, fat and protein to produce energy and affect enzymes that influence the muscles, nerves, and heart.
- Vitamin B₂ (Riboflavin) helps the body process amino acids and fats, makes antibodies and red blood cells and maintain good vision & feathers.
- Vitamin B₆ (Pyridoxine) helps the body break down protein, and helps maintain the health of red blood cells, the nervous system, and parts of the immune system.

Prithi[®] WS powder, is specially used in the treatment of various manifestations of its component vitamin deficiency such as polyneuritis, curled toe paralysis, inappetance.

Prithi[®] WS powder, supplementary vitamins preparation for poultry.

Indication:

Prithi[®] WS Powder is specially used in the treatment of various manifestations of its component vitamin deficiency such as Polyneuritis, Curled toe paralysis, inappetance or where requirement are increased (carbohydrate-rich diet, pregnancy).

Dosage & Administration:

Poultry: 1 gm / 1 litre drinking water or, 200 gm / 100 kg feed for 3-5 days.

Contraindication & Precaution:

Prithi[®] WS Powder is contraindicated in birds and animals hypersensitive to any of its components.

Side effect:

It has no side effect at recommended dose.

Side effects have been reported with specific vitamins & minerals with higher dosage.

Storage Condition:

Store in a cool & dry place, out of reach of children. Protect from light and moisture and keep away from feed. Use the content as early as possible after opening. Keep away out of reach of children, protect from light & moisture.

How Supplied:

100 gm in plastic container.