

Sulprim-Vet[®] Suspension

Sulphadiazine BP & Trimethoprim BP

Composition:

Each ml suspension contains Sulphadiazine BP 400 mg & Trimethoprim BP 80 mg.

Indication:

Sulprim-Vet[®] suspension is used in respiratory, gastrointestinal, urinary & reproductive infections in poultry. It is used for the treatment and prevention of different infections caused by gram positive bacteria, gram negative bacteria and mycoplasma in poultry e.g. colibacillosis, salmonellosis or pullorum disease, fowl typhoid, pasteurellosis or fowl cholera, gastroenteritis, chronic respiratory infection (CRD) & secondary infection of viral diseases. Sulprim-Vet[®] suspension is also used when other antibiotic and chemotherapeutic agents are not effective due to development of resistance.

Dosage and Administration:

Sulprim-Vet[®] suspension should be used daily with drinking water. At first required dose should be mixed with little quantity of water which is then mixed with total quantity of water for easy dissolution.

Poultry treatment: 1 ml. suspension with 5 litres of drinking water daily for 4-6 days.

Poultry prevention: 1 ml suspension with 10 litres of drinking water daily for 2-3 days.

Contraindication:

The drug is contraindicated in animals hypersensitive to active ingredients.

Withdrawal Period:

Poultry (Broiler) treated with this drug should not be slaughtered for human consumption for a period of 5 days after last treatment

Side Effect:

At the recommended dose, side effects are rare and mild.

Precaution:

Shake the bottle before use.

Store in a cool, dry place. Protect from light and keep out of reach of children. To be dispensed only by or on the prescription of a registered veterinary doctor.

Pack Size:

100 ml suspension.