

Sumid-Vet[®] Powder

Sulphanilamide Powder BP

Composition:

Sumid-Vet[®] Powder contains cream-white, odourless, crystalline Sulphanilamide powder BP 100%.

Indication:

- ✓ Sumid-Vet[®] Powder is used to treat sulphanilamide -sensitive organisms, notably streptococcal septicaemia; equine pneumonia, strangles, joint-ill, metritis in mares, cows and ewes.
- ✓ Sumid-Vet[®] Powder is also used as a dusting powder for wounds by topical application.

Dosage and Administration:

For Oral Administration:

Animals up to 3 months of age:

Initial dose: 7.5 gm per 50 kg body weight.

Continuation: 2.5 gm per 50 kg body weight every eight hours interval.

Older animals:

Initial dose: 10 gm per 50 kg body weight.

Continuation: 5 gm per 50 kg body weight every twelve hours interval.

Total period of treatment should not normally exceed 7 days.

Or as directed by the registered Veterinary physician.

For external use:

Use externally as per requirement or prescribed by a registered veterinary doctor.

Contraindication:

In therapeutic doses Sumid-Vet[®] Powder is relatively non-toxic, but prolonged treatment has been known to cause agranulocytosis, haemolytic anaemia and avitaminosis-K. Where prolonged treatment is undertaken, patients should be watched with care. Sumid-Vet[®] Powder occasionally causes crystalluria, particularly when urinary pH is low. Always ensure adequate water intake during treatment and take particular care in the case of animals suffering from renal damage.

Side Effect:

Includes allergic rashes, diarrhoea, dehydration, debility, loss of appetite, hypogalactia, digestive disturbances in ruminants and in other species depression, nausea, vomiting, crystalluria, haemolytic anaemia.

Storage:

Protect from light, store in a cool and dry place. Keep out of reach of children.

Pack Size:

10X10 gm.