

Myonil[®]

Eperisone Hydrochloride

COMPOSITION

Myonil[®] Tablet: Each sugar coated tablet contains 50 mg of Eperisone Hydrochloride INN.

INDICATIONS

Improvement of muscular hypertonic symptoms in the following diseases Cervical syndrome, periarthrititis of the shoulder, lumbago. Spastic paralysis in the following disease: Cerebrovascular disease, spastic spinal paralysis, cervical spondylosis, postoperative sequelae (including cerebrospinal tumor), sequelae to trauma (spinal trauma, head injury), amyotrophic lateral sclerosis, cerebral palsy, spinocerebellar degeneration, spinal vascular diseases and other encephalomyelopathies.

DOSAGE AND ADMINISTRATION

For adults:

Usually 3 tablets per day in three divided doses after each meal. The dosage should be adjusted depending on the patient age and severity of symptoms.

CONTRAINDICATION

Myonil is contraindicated in patients with a history of hypersensitivity to Eperisone Hydrochloride.

USE IN PREGNANCY AND LACTATION

Eperisone should only be used in pregnant women if the expected therapeutic benefits are evaluated to outweigh the possible risks of treatment. The safety of Eperisone has not been established in pregnant women. The drug should not be used during lactation.

SIDE EFFECTS

The side effects of Eperisone are very rare, only a few cases have been observed. These are excessive relaxation, stomachache, nausea, vertigo, anorexia, drowsiness, skin rashes, diarrhea, vomiting, indigestion, GI disturbances, insomnia, headache, constipation, etc.

STORAGE

Store in a cool and dry place, protect from light and moisture. Keep all medication out of reach of children.

HOW SUPPLIED

Myonil[®] Tablet: Each box contains 5x10's in blister packing.

Manufactured by



SQUARE
PHARMACEUTICALS LTD.
BANGLADESH