

Iverzol-Vet[®] Suspension

Ivermectin BP & Triclabendazole INN

Composition:

Each ml Iverzol-Vet[®] Suspension contains Ivermectin BP 2 mg & Triclabendazole INN 120 mg.

Indication: Iverzol-Vet[®] Suspension is indicated for the control of adult and immature (including inhibited stages) ivermectin-sensitive gastrointestinal nematodes such as *Ostertagia sp.*, *Haemonchus sp.*, *Trichostrongylus sp.*, *Cooperia sp.*, *Oesophagostomum sp.*, *Nematodirus sp.*, *Strongyloides sp.*, *Chabertia sp.*, *Trichuris sp.*, lung worms (*Dictyocaulus sp.*) in cattle and sheep. As well as sucking lice in cattle (*Linognathus sp.*, *Haematopinus sp.*, *Solenopotes sp.*), Itch mite (*Psorergates sp.*) and nasal bots (all larval stages of *Oestrus sp.*) in sheep and strains of above parasites resistant to other classes of parasiticides. Iverzol-Vet[®] Suspension is also indicated for adult, immature and early immature triclabendazole-sensitive liver flukes (*Fasciola sp.*) in cattle and sheep.

Dosage & Administration: The recommended dose rate of Iverzol-Vet[®] Suspension: **Cattle:** 5ml per 50 kg bodyweight, corresponding to 0.2 mg ivermectin and 12 mg triclabendazole per kg. It is given orally. Should shake well before use. It is suitable for use through most types of drenching guns. *Treatment of sick animals suffering from worm infections:* When sick animals are treated all other animals with the same history should be treated as well. Possible subsequent administrations should be given according to veterinary advice.

Withdrawal Period: Milk: 28 days, Meat: 21 days.

Precaution: Keep out of reach of children.

Storage: Store in a dry and cool place. Protect from light. Keep out of reach of children.

Pack size: 50 ml.

For veterinary use only. To be dispensed only on the prescription of a registered veterinary doctor.