

Floricol-Vet Solution

Florfenicol INN 20%

Composition

Each ml solution contains Florfenicol INN 200 mg.

Indication

Floricol-Vet™ solution is indicated for preventive and therapeutic treatment of gastrointestinal and respiratory tract infections, caused by Florfenicol sensitive micro-organisms such as *Actinobaccillus* spp., *Pasteurella* spp., *Salmonella* spp. and *Streptococcus* spp. in animal and poultry. The presence of the disease in the herd should be established before preventive treatment. Medication should be initiated promptly when respiratory disease is diagnosed.

Pharmacology

Florfenicol is a synthetic broad-spectrum antibiotic that is effective against most Gram-positive and Gram negative bacteria isolated from domestic animals. Florfenicol, a fluorinated derivative of chloramphenicol, acts by binding to the 50s ribosomal subunit of bacteria, thereby inhibiting bacterial protein synthesis and is bacteriostatic.

Dosage & Administration

1 ml /2 L drinking water (20-30 mg/Kg body weight) for 3-5 days. Stir continuously while mixing with water. Or, as per direction of a registered Veterinarian/Consultant.

Contra-indication

Do not administer in case of previous hypersensitivity to Florfenicol.

Side-effects

A decrease in food and water consumption and transient softening of the faeces or diarrhoea may occur during the treatment period. The treated animals recover quickly and completely upon termination of treatment.

Precautions

Can be used only under supervision of a registered Veterinarian. Complete full course according to the prescription. Dosage and administration must be as per prescription

Withdrawal Period

Broiler: 5 days.

Storage

Store below 30°C temperature, protected from light. Keep out of reach of children.

Pack Size

100 ml HDPE bottle, 500 ml HDPE bottle.