

Ampicin-Vet[®] Injection

Ampicillin Sodium BP

Composition:

Each vial contains Ampicillin Sodium BP 2.12 gm equivalent to 2.0 gm Ampicillin along with 10 ml WFI.

Indication:

Indicated in different bacterial diseases of animals e.g. Pneumonia, Foot rot, Enteritis, Salmonellosis, Calf scour, Infected wound, Fever, Colibacillosis, Metritis, Mastitis, Pyelonephritis, Black quarter, Joint infection, Kidney infection, Haemorrhagic septicemia etc.

Highlighting Points:

- True broad spectrum antibiotic.
- Low resistance in veterinary use.
- Well absorbed following IM or IV administration.
- Rapidly and widely distributed into most body fluid.
- Ensures quicker action.
- Safe in pregnancy and lactating period.
- Reduces the chance of diarrhea by parenteral administration
- The half life is approximately 60-90 minutes.
- Tissue drug concentrations may be higher than serum concentrations.

Dosage & Administration:

Ampicin-Vet[®] powder for injection should be mixed with 10 ml WFI and then administer by IM or IV route. Parenteral dose is 7 mg/kg body wt.

Animals	Body Wt.	Dose	Dosage Interval
Cattle, Buffalo	50 Kg	1.0 to 2.5 ml	Once daily but in severe infection twice daily.
	100 Kg	2.0 to 5.0 ml	
	200 Kg	4.0 to 10.0 ml	
Goat, Sheep	10 Kg	0.2 to 0.5 ml	
	20 Kg	0.4 to 1.0 ml	
Cat, Dog	10 Kg	0.5 to 1.0 ml	
	20 Kg	1.0 to 2.0 ml	

Contraindication:

The drug is contraindicated in animals hypersensitive to Ampicillin.

Precaution:

For veterinary use only. To be dispensed only by or on the prescription of a registered veterinary doctor. Store in a cool, dry place. Protect from light and keep out of reach of children.

Side Effects:

In recommended dosage, Ampicin-Vet[®] is relatively non-toxic and free from adverse effect. Moreover, skin rash, pain at the site of injection, diarrhea etc. may occur.

Use in Pregnancy & Lactation:

Ampicin-Vet[®] is safe in pregnancy state and in lactating period in animals.

Storage Condition:

Store in a cool & dry place. Protect from sunlight & moisture. Keep out of reach of children.

Pack Size:

Each pack encloses a vial containing Ampicin-Vet[®] PFI with an ampoule of 10 ml WFI.