

# Cotrim-Vet<sup>®</sup> Bolus

Sulphamethoxazole BP & Trimethoprim BP

**Composition:**

Each bolus contains Sulphamethoxazole BP 1000 mg & Trimethoprim BP 200 mg.

**Indication:**

It is used for infections of urinary, respiratory and genital tracts, skin and wound infections and other bacterial infections. It is used for the treatment of actinobacillosis, actinomycosis, coccidiosis, mastitis, canine ehrlichiosis, oral necrobacillosis, infectious foot rot, necrotic rhinitis, metritis, infectious keratitis, infectious polyarthritis, post abortion, pneumonia, salmonellosis, toxoplasmosis, bronchitis, diarrhoea, bronchopneumonia, etc.

**Dosage and Administration:**

Cattle & Horse : 30 mg/kg body weight or 1 bolus/40-kg body weight daily.

Calf, Sheep, Goats : 30mg/kg body weight daily for 5 to 7 days.

Cotrim-Vet<sup>®</sup> can be administered orally by mixing with foods or drinking water.

**Use in Pregnancy and Lactation:**

Risk of haemolysis & kernicterus in calves treated with this drug.

**Contraindication:**

Cotrim-Vet<sup>®</sup> should not be administered to ruminants with functioning rumen to avoid possible disturbance of the ruminal flora. It is also contraindicated in animal with severe hepatic or renal insufficiency, blood disorders.

**Side Effect:**

Frothing and salivation can follow oral dosing in cats. Folate deficiency may occur during prolonged treatment in animals.

**Precaution:**

For animal use only. Adequate fluid intake should be ensured during therapy.

**Package Quantity:**

5 x 4's blisters in a box.