

HamecoMoltox-Dry[®] Powder

Mold inhibitor, Toxinbinder, Mycotoxin bio-synthesis inhibitor & Mycotoxin bio-nutralizer

Composition:

HamecoMoltox-Dry[®] contains – Propionic acid, Calcium propionet, Fumeric acid, Sorbic acid, and as active career Hydrated Magnesium Aluminosilicate.

Uses:

- Acts as a Toxin binder and destroy Mycotoxin from poultry feed.
- Acts as a broad spectrum Mould inhibitor & destroy fungus from poultry feed.
- It prevents bio-synthesis of mycotoxin in feed.
- It causes bio-nutralization of Mycotoxin when feed contain Mycotoxin.
- It enhances growth rate of poultry.
- It increases egg production in layer.
- Absorb moisture from feed & keep feed in dry condition.
- It promotes enzymatic activity & liver function for detoxification.

Dosage Rate:

100-200gm / 100 kg feed continue. Or, 250 gm / 100 kg feed when feed is affected severely with Fungus, Mould & Mycotoxin.

To get better result keep the mixture feed 24 – 48 hours before feeding.

Side Effect:

It has no side effect

Stability:

It remains stable up to 2 years when store in a cool & dry place.

Precaution:

Store in a cool & dry place. Use as early as possible after opening the container. Protect from direct sunlight & moisture. Keep away from reach of children.

Packing:

25 kg bag.