

Otetra-Vet[®] LA Injection

Oxytetracycline BP
Long Acting Injectable Solution

Composition:

Each ml Otetra-Vet[®] LA injection contains Oxytetracycline 200 mg as Oxytetracycline Dihydrate BP.

Indication:

For the treatment and prevention of different types of bacterial diseases of cattle, buffalo, sheep, goat, deer, rabbit and poultry.

Cattle, Buffaloes, Sheep, Goat & Deer: For the treatment of actinobacillosis, bacterial enteritis, bacterial pneumonia, bovine respiratory disease complex, diphtheria, keratoconjunctivitis (pink eye), leptospirosis, metritis, anaplasmosis, foot-rot, black leg, navel ill, joint ill, heart water, dermatophilosis, enzootic abortion (bedsoniasis), wound infections, mycoplasmosis, Aid in the control or prevention of shipping fever, pneumonia, post operative and post parturient infections.

Rabbit: For the prevention and treatment of conditions / diseases susceptible to Oxytetracycline including infections due to Pasteurella multocida, Bordetella bronchiseptica, Escherichia coli and Salmonella spp etc.

Poultry: For the prevention and treatment of conditions / diseases susceptible to Oxytetracycline including fowl cholera, fowl typhoid, complicated CRD etc.

Benefits of Otetra-Vet[®] LA Injection:

- Otetra-Vet[®] LA Injection is a highly active broad spectrum antibiotic which gives long time effects after administration.
- Single dose of Otetra-Vet[®] LA Injection is sufficient to ensure full treatment.
- Otetra-Vet[®] LA Injection ensures long time (3-5 days) effective plasma concentration after single dose administration. Therefore no repeated dose is required.
- Otetra-Vet[®] LA Injection prevents growth of different bacteria, protozoa & intercellular microorganisms like mycoplasma, chlamydia, rickettsia etc.
- Otetra-Vet[®] LA Injection is used for the treatment & prevention of diseases caused by gram positive & gram negative bacteria in cattle, buffalo, sheep, goat, deer, rabbit, & poultry.

Dosage and Administration:

Cattle, Buffaloes, Sheep, Goat & Deer: Single dose treatment of 1 ml per 10 kg body weight. (or, 20 mg per kg body weight) by IM route.

Poultry & Rabbit: Single dose treatment of 0.25 ml per kg body weight (or, 50 mg per kg body weight) by subcutaneous route.

Severe infections may require re-treatment at 3 to 5 days following initial injection.

Or, as advised by the veterinary physician.

Contraindication:

Don't administer those animals which are hypersensitive to Oxytetracycline. The use of Oxytetracycline should be avoided in animals with an impaired liver and / or kidney function.

Precaution:

To minimize local irritation at the site of injection in cattle, it is recommended that not more than 10 ml should be injected at any one site.

Withdrawal Period: Meat - 28 days & milk - 4 days.

Storage:

Store below 30° C temperature. Protect from light. Keep out of reach of children.

Package quantities:

10 ml vial