

# Stress-Kill<sup>®</sup> WS Powder

Water Soluble Multivitamin

## Composition:

Each 100 gm powder contains -

Vitamin A	25,00,000	IU	Folic Acid	125.00	mg
Vitamin D <sub>3</sub>	3,00,000	IU	Biotin	10.00	mg
Vitamin E (50%)	5,000.00	mg	Cobalt Sulphate, 7H <sub>2</sub> O	10.00	mg
Vitamin K <sub>3</sub> (10%)	2,000.00	mg	Copper Sulphate, 5H <sub>2</sub> O	150.00	mg
Vitamin B <sub>1</sub>	1,250.00	mg	Ferrous Sulphate, 7H <sub>2</sub> O	100.00	mg
Vitamin B <sub>2</sub>	2,500.00	mg	Potassium Iodide	10.00	mg
Vitamin B <sub>6</sub>	1,250.00	mg	Manganese Sulphate, 1H <sub>2</sub> O	1,000.00	mg
Vitamin B <sub>12</sub> (0.1%)	250.00	mg	Zinc Sulphate, 7H <sub>2</sub> O	1,000.00	mg
Nicotinamide	2,500.00	mg	DL Methionine	1,000.00	mg
Calcium-D Pantothenate	2,500.00	mg	L-Lysine	2,000.00	mg
Vitamin C	5,000.00	mg			

## Benefits:

- Increases growth & improves meat quality of broiler.
- It increases egg production, fertility & hatchability.
- It prevents faulty growing & weakness of bone.
- It increases favorable bacteria in intestine & helps in adequate Vitamin-B complex formation.
- In large animal Stress-Kill increases growth rate of calf & production capacity of adult.
- In case of shifting, vaccination, de-worming, treatment of parasitic & bacterial diseases Stress-Kill gives better result & prevents weakness.

## Indication:

Stress-Kill<sup>®</sup> is indicated for the treatment and / or prevention of vitamin & mineral deficiencies associated with poor feeding, improper feed, and lower feed intake and decreased absorption. It is also indicated in birds with increased requirements for vitamins & minerals due to acute and chronic diseases, infections, during treatment with antibiotics / anthelmintics etc.; and any kind of stress, e.g. - Weakness after birth, adverse change of temperature, replacement, change of feed formulation, transportation & Vaccination.

## Dosage & Administration:

Poultry: 1 gm / 5 litre of drinking water or, 20-40 gm / 100 kg feed for 4-5 days.

## How Supplied:

100 gm sachet.