

Two-Plus Liquid

Liquid toxin binder for poultry

Composition: Each litre contains

Yeast extracts	0.30 gm	Phosphoric acid	6.50 gm
Citric acid	8.00 gm	Malic acid	1.00 gm
Tartaric acid	1.50 gm	Riboflavin	0.30 gm
Na K tartrate	0.10 gm	Na citrate	0.10 gm
Thiamin Mononitrate	0.30 gm	Pyridoxine HCl	2.00 gm
Cu Penta-sulphate	10.00 gm	Di-sodium EDTA	1.50 gm
Propylene glycol	100.00 gm	Aspartic acid	0.20 gm
Sacharomyces cerevisae	20.00 gm	Distilled water	q.s. to make 1 litre

Indication:

Overcome mycotoxicosis, reduces the negative effect of mycotoxins, improves response to vaccination, improves animal immune system, better functioning of body organs especially liver kidney and ovaries and improves amino acid absorption.

Dosage & Administration:

1ml/2-4 liter of drinking water for 3-5 days.

Side Effect:

It has no side effect at recommended dose.

Drug Interaction:

Two-plus does not interfere with the activity of other antibiotics in case of Mycotoxicosis.

Storage condition:

Store in a cool & dry place. Protect from sunlight & moisture. Use as early as possible after opening. Keep away from reach of children.

Pack size: 100ml, 500 ml & 1litre.